

# InTice<sup>TM</sup> Select ant granules

Kills Ants, Roaches, Crickets, Mole Crickets, Silverfish, Sowbugs, Pillbugs, Earwigs, Centipedes, and Millipedes

**Weather/Moisture Resistant Bait  
For Indoor and Outdoor Use  
Ready-to-Use Bait**

Active Ingredient: Orthoboric Acid.....5%	EPA Reg. No.: 73079-2
Other Ingredients:.....95%	EPA Est. No.: 73079-MO-1
Total:.....100%	Net Contents: 1 lb

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### Use Restrictions

In food areas of restaurants, residential, health care, educational, daycare, agricultural buildings, animal rearing and handling establishments, food handling and food processing establishments, this product may only be applied as a crack and crevice treatment. Any product remaining outside of cracks and crevices must be cleaned up and removed.

### Application Directions

#### Sites and Species

**Use Sites:** In and around residential, multi-family, commercial, industrial, municipal, institutional, research, recreational, health care, educational, daycare, hospitality and agricultural buildings and other man-made structures, garages, transport vehicles, sewers, animal rearing and handling establishments and food handling and food processing establishments. Turf sites include lawns, ornamental turf, parks, playing fields, right-of-ways, and golf courses.

**Listed Species: Ants:** Odorous House, Argentine, Carpenter, Pavement, Little Black, Pharaoh, Crazy, Acrobat, Rover, Big-Headed, White-Footed, Ghost, Cornfield, and Thief; **Roaches:** German, American, Brown-Banded, Smokey Brown, Brown, Asian, Wood and Oriental; **Crickets:** House, Field, Camel, Ground, and Mole; **Others:** Silverfish, Earwigs, Sowbugs, Pillbugs, Centipedes and Millipedes.

**Outdoor Perimeter Application:** Apply bait at a rate of 4 lbs per 1000 square feet (6 oz or 1.25 cups per 100 square feet). Use a hand shaker, duster or mechanical spreader to apply the bait. Apply in a band from two up to four feet wide around the perimeter of buildings. Also apply around exterior light fixtures and along the edges of sidewalks, patios and driveways and under decks. Pay particular attention to potential insect harborage sites such as mulch beds, flowerbeds, compost heaps, wood piles, trees, stumps and trash areas and apply bait at the more liberal application rate in these areas. Do not apply the bait directly to paved or similar hard surfaces and do not pile the bait. The bait may also be sprinkled in and around rodent bait stations to control insects that harbor in these stations. When treating for ants, apply 1/2 to 1 oz of bait around ant mounds and sprinkle along foraging trails.

## KEEP OUT OF REACH OF CHILDREN CAUTION

### PRECAUTIONARY STATEMENTS

**Hazards to Humans and Domestic Animals: CAUTION:** Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. **Personal Protective Equipment (PPE)** Applicators and handlers must wear: Long-sleeved shirt and long pants: chemical resistant or waterproof gloves: and shoes plus socks.

#### First Aid

**If in Eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first five minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

**If on Skin:** Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

Have the product or label with you when calling a Poison Control Center or doctor or going for treatment. In case of emergency, for additional information call **800 858 7378**.

**Environmental Hazards:** Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

For ants, including Carpenter ants, pay particular attention to harborage areas and apply bait around trees and stumps, in tree cavities, in and around firewood piles, around landscaping stones and in mulch beds. Ants can take the bait back to the nest and kill the colony. Bait may also be applied in insect bait stations. Reapply bait after one week if infestation is still present. To prevent large insect populations from developing around structures in warm months, a quarterly application around the perimeter and to harborage areas, as described above, at a rate of 1 lb per 1000 square feet is recommended. This treatment is particularly effective to help prevent the establishment of large colonies of pest ants. InTice Granular Bait may be used in combination with registered non-repellent insecticides (products containing the active ingredients imidacloprid or fipronil). If the perimeter is treated with a repellent insecticide (eg. pyrethroid), the bait may still be used around the perimeter; but it should be applied in a band outside the repellent barrier for best results. As a separate matter from the use of this product, as part of an IPM program, it is recommended that overhanging trees and other vegetation touching the structure be trimmed back, and cracks or gaps around pipes and window and door frames be sealed to minimize potential insect entry into buildings.

**Sewer Application:** Apply bait along ledges inside sewers and around manhole covers.

**Indoor Application:** Apply bait only in areas inaccessible to children and pets, or in tamper-resistant bait stations. Apply bait at a rate of 6 oz (1.25 cups) per 100 square feet. Using a hand shaker or duster, apply bait in attics, crawl spaces, trash closets, utility closets, dropped ceilings, on dirt or gravel cellar floors, under and behind appliances and vending machines, in wall voids and voids under and behind cabinets, sinks and tubs, in drawer wells, garbage chutes, pipe chases, elevator pits, around drains, around electrical conduits and in cracks and crevices where insects may harbor. Focus application in areas where insects or their signs are seen. Reapply bait after one week if infestation is still present. A power duster may be used to apply bait in appropriate areas, such as attics and crawlspaces. The duster manufacturer's recommendations for settings should be followed.



toll free: 866 788 4101  
www.rockwelllabs.com