



# MAXFORCE<sup>®</sup>

## Fly Spot Bait

<b>ACTIVE INGREDIENT:</b> Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]-N nitro-2 imidazolidinimine	10.0%
2,9-Tricosane	0.1%
<b>OTHER INGREDIENTS:</b>	89.9%
<b>TOTAL:</b>	100.0%
<b>EPA Reg. No. 101563-119</b>	<b>EPA Est. No.</b>

**STOP - Read the label before use.**  
**Keep Out of Reach of Children.**  
**CAUTION**

See Panel for First Aid Instructions and for Complete Precautionary Statements and Directions for Use.  
 For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-424-9300  
 For **PRODUCT USE** Information Call 1-800-331-2867

FIRST AID		PRECAUTIONARY STATEMENTS	
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>	<b>HAZARDS TO HUMANS AND DOMESTIC ANIMALS</b>	<p><b>CAUTION:</b> Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Wear long-sleeved shirt and long pants, socks, and shoes.</p>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>	<b>ENVIRONMENTAL HAZARDS</b>	<p>This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.</p>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>	<b>PRODUCT FACTS</b>	
<p>In case of emergency call toll free the Environmental Science U.S., LLC Emergency Response Telephone No. 1-800-424-9300. Have the product container or label with you when calling a poison control center or doctor or going for treatment.</p> <p><b>Note To Physician:</b> No specific antidote is available. Treat the patient symptomatically.</p>			
<p>• Contains a specially formulated mixture of attractants. • Controls house flies resistant to organophosphates, carbamates, and pyrethroid insecticides. This product also controls *fruit flies. Applications of this product can aid in suppression of psychodid and phorid flies.</p>			
<b>MAXFORCE<sup>®</sup> FLY SPOT BAIT</b>	<ul style="list-style-type: none"> <li>A bait that is consumed by listed flies.</li> <li>Easy to apply</li> <li>Outdoor residual efficacy up to 14 days unless the bait is completely consumed prior.</li> <li>In animal facilities and protected areas, residual efficacy can last up to 6 weeks unless the bait is completely consumed.</li> </ul>	<ul style="list-style-type: none"> <li>Contains the attractant muscature</li> <li>Contains 2 powerful attractants designed to lure listed flies to their death.</li> <li>Contains Bitrex<sup>®</sup> bittering agent.</li> </ul>	
<b>WHERE TO APPLY</b>	<ul style="list-style-type: none"> <li>In and around agricultural livestock production facilities, including: poultry houses, turkey and chicken houses, cattle, sheep and goat facilities, feedlots, dairy barns, swine confinement buildings, dog kennels, and horse stables.</li> <li>Around the outside of commercial and residential structures, or partly enclosed or protected areas</li> </ul>	<ul style="list-style-type: none"> <li>In non-food/feed areas of commercial facilities</li> <li>*Applications for fruit fly control may only be made indoors in California.</li> </ul>	
<b>HOW TO APPLY</b>	<ul style="list-style-type: none"> <li>Mix in water and spray or paint on surfaces.</li> </ul>		
<b>COVERAGE</b>	<ul style="list-style-type: none"> <li>Use 16 oz of product in 1 gallon of finished spray to treat surfaces where listed flies may rest in buildings with 2,000 sq ft of floor space.</li> </ul>		
<b>RE-APPLY</b>	<ul style="list-style-type: none"> <li>Re-apply as needed, but not before 7 days.</li> </ul>		
<b>QUESTIONS ??</b>	<ul style="list-style-type: none"> <li>For questions or comments, call toll-free 1-800-331-2867</li> </ul>		

**DIRECTIONS FOR USE**  
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling.  
**IMPORTANT: Read entire label before using this product.**

D00000956  
 81779600F 221216AV1  
**Net Weight: 2 oz (56.7 g)**



PRODUCED FOR  
 Environmental Science U.S., LLC  
 5000 CentreGreen Way, Suite 400  
 Cary, NC 27513  
 Product of Austria

**Best Management Practices**

- Use the bait as part of an overall Integrated Pest Management program.
- Use proper sanitation and manure management.
- Use environmental controls, such as proper light management.
- Consult with cooperative extension service in your state for advice on listed fly management practices.

**Product Information**  
 This product is a bait formulated to control listed flies in rural and urban environments, in animal confinement areas (e.g., cattle, poultry, swine), in commercial establishments, and outdoors in residential areas. The product can be used as a spray for treatment of surfaces, such as stable walls or waste containers, or it can be used as a point-on application to walls and other surfaces such as pillars and compartment walls. There is no need to spray the complete surface because this bait will attract pests to the treated area. Begin applications at the start of the season before listed fly populations have reached their peak. Apply the product to many locations throughout the structure or facility. Spray the product on surfaces where listed flies tend to rest. Begin applications at the start of the season before listed fly populations have reached their peak.

**How and where to apply**

**FOR USE IN** broiler houses, turkey houses, caged layer houses, feedlots, dairy barns, swine confinement buildings, horse stables, kennels, and other animal confinement facilities where listed flies tend to be a nuisance or a health hazard.

**Spray application**  
 Spray diluted product where listed flies aggregate: on walls, around windows, on surfaces of pillars and other structures, etc. Apply product in areas of 3 to 12 sq ft where listed flies tend to rest. If treating areas 10 sq. ft. in size, you can apply up to 50 such bait areas in a building with 1,000 sq ft of floor space without exceeding the limit of 8 ounces of product for a building of this size.

**Paint on application.**  
 Paint the product in small bands up to 6 inches wide, or more, on places where listed flies aggregate. Apply this product at a rate of 8 ounces for every 1,000 sq ft of floor space.

**Residual efficacy in animal facilities.**  
 Residual efficacy in animal facilities is dependent on the sanitary conditions and listed fly pressure level, as well as on the quality and cleanliness of the surfaces treated.

Apply as soon as listed fly pressure exceeds acceptable threshold levels to avoid population build up. Re-apply as needed, typically at 4 to 6 week intervals.

When spraying surfaces, care must be taken only to spray places out of reach of the animals. Do not spray directly on animals, feed and drinking supplies nor on milking equipment. This also applies to empty pens due to be re-stocked within the next few weeks. Do not treat surfaces that people frequently contact. Avoid contact with sprayed surfaces until dry.

**FOR USE IN OTHER AREAS** such as non-food/non-feed areas of commercial food handling establishments, restaurants, and dining areas, residential backyards, on and around dumpsters and trash collection areas, storage facilities, dog kennels, zoos, animal rearing facilities, schools, hospitals, nursing homes, offices, marinas, and other locations where listed flies tend to aggregate and are a nuisance or a health hazard.

**Spray application**  
 Spray diluted product where listed flies aggregate: on walls, around windows, on surfaces of dumpsters or trash collection bins, etc. For best results, apply product on areas of 3 to 12 sq ft where listed flies tend to rest. If treating areas 10 sq ft in size, you can apply up to 50 such bait areas in a building with 1,000 sq ft of floor space without exceeding the limit of 8 ounces of product for a building of this size.

**Paint on applications**  
 Paint the product in small bands up to 6 inches wide, or more, on places where listed flies aggregate. Apply this product at a rate of 8 ounces for every 1,000 sq ft of floor space.

**Food/Feed Handling Establishments**  
 Applications in food/feed handling establishments are permitted ONLY in non-food/non-feed areas. Non-food/non-feed areas include garbage rooms, lavatories, entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, mop closets and storage (after packaging, canning or bottling). Food/feed handling establishments are defined as places other than private residences in which exposed food/feed is held, processed, prepared or served. Included also are areas for receiving, storing, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups) of food.

Serving areas, when food is exposed and the facility is in operation, are also considered food/feed areas. However, applications of this product may be made in serving areas ONLY while the facility is not in operation, and provided exposed food has been covered, or removed from the area being treated prior to application. These "spot" applications may cover no more than 2 sq ft per treated area, may not be spaced closer than 10 ft apart, and should be a minimum of 6 feet above floor level.

Contamination of food or food contact surfaces must be avoided. Do not apply this product to any food, food packaging, or any food contact surfaces.

Applications of this product may leave a visible residue or discolor some surfaces. Before treating, it is advisable to test application in a small inconspicuous area.

**Mixing Directions and Use Rates**  
 For spray applications, add the contents of one 2 ounce packet of this product to 1 pint of water and agitate until completely mixed. One pint of finished spray will treat 125 sq ft of surfaces where listed flies may rest. For paint-on applications, add 8 ounces of this product to 8 ounces of water (approximately 1 cup), and mix to desired consistency for results. Apply this mixture in narrow bands up to 6 inches wide on suitable surfaces where listed flies may rest. Do not apply this product at a rate exceeding 8 ounces for every 1,000 sq ft of floor space.

**Application equipment**  
 For spray applications, apply as a coarse fan or cone spray. Apply solution at a low pressure not to exceed 20 psi. Flush spray equipment with water to prevent clogging.

For paint-on applications, use a brush or roller paint applicator to apply solution to surfaces. Clean application and mixing equipment with water after application.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.  
**Pesticide Storage:**  
 Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, or feed. Store in original containers and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage.

**Pesticide Disposal:**  
 Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Handling:**  
 For less than one pound size. Non-refillable container. Do not reuse or refill this container. Completely empty container. Wrap used container and place in trash can.

**Spills:**  
 If the container is leaking or material is spilled, carefully sweep up spilled bait and wash exposed area. In spill or leak incidents, keep unauthorized people away.

You may contact the Environmental Science U.S., LLC Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Environmental Science U.S., LLC Emergency Response Telephone No. is 1-800-424-9300.

### IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Environmental Science U.S., LLC. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ENVIRONMENTAL SCIENCE U.S., LLC MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Environmental Science U.S., LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ENVIRONMENTAL SCIENCE U.S., LLC DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT ENVIRONMENTAL SCIENCE U.S., LLC'S ELECTION, THE REPLACEMENT OF PRODUCT.

Envu and the Envu logo are trademarks and MAXFORCE<sup>®</sup> is a registered trademark owned by Environmental Science U.S., LLC or one of its affiliates. Bitrex<sup>®</sup> and Contains Bitrex<sup>®</sup> are registered trademarks of Johnson Matthey Public Limited Company.



- PMS 663
- PMS 2383
- PMS 2081
- Black

**NOTE: Full eyespot may alternatively be located in lower left corner of Back Panel**